MIDI PARASEP®

Faecal parasite concentrator helminth ova and larvae/protozoa cysts and oocysts





APACOR

MIDI PARASEP®

Faecal parasite concentrator

Health & safety benefits

- Totally enclosed/sealed process
- Reduced reagent volumes
- No cleaning required
- Single use, no sample contamination
- Ready to use systems available

Performance benefits

- Optimum sample recovery
- Enhanced sample clarity
- Rapid four step process
- Human resources optimised
- Easy patient identification
- Fits all 50ml centrifuge buckets

Step 1

Code

145000

145002

145003

145300

145400

145501

Sample preparation Introduce sample to the fixative/solvent

Step 2 **Emulsification** Vortex or shake

2

Midi Parasep - 50ml Intestinal Parasite Concentrators:

6ml Apafix™ + 80ml Ethyl Acetate

6ml Formalin & Triton X + 80ml

6ml SAF & Triton X + 80ml

6ml Formalin & Triton X

to emulsify



3

Debris

Fatty plug supernatant Sediment

UOM

ΡK

ΡK

ΡK

ΡK

ΡK

ΡK

Pack size

50

40

40

40

40

40

Self standing sample chamber

Mixing chamber

Integral spoon


425µm Filter

Debris trap


Air/liquid seal & safety lock

Sedimentation cone

Step 4 Examination Slide preparation



Note: To calculate the required RPM for any centrifuge.



RPM - rotor speed in

- revs/min. g - centrifugal force
- (max. 1000g) - radius, horizontal
- distance between sedimentation cone
- tip and spindle centre measured in mm.

APACOF

Visit www.apacor.com for all our latest information or contact us at sales@apacor.com

www.apacor.com APA-BR-015/06

Description

No Additive

6ml Apafix™

Ethyl Acetate

Ethyl Acetate

SAPPHIRE CENTRE FISHPONDS ROAD, WOKINGHAM BERKSHIRE, RG41 2QL, ENGLAND TEL: +44 (0)118 979 5566 FAX: +44 (0)118 979 5186